

Part VI: Summary of the Risk Management Plan

Summary of Risk Management Plan for Teriflunomid-ratiopharm 14 mg Filmtabletten

This is a summary of the risk management plan (RMP) for Teriflunomide-ratiopharm 14 mg film-coated tablets (hereinafter referred to as Teriflunomide). The RMP details important risks of Teriflunomide, how these risks can be minimised, and how more information will be obtained about Teriflunomide risks and uncertainties (missing information).

Teriflunomide summary of product characteristics (SmPC) and its package leaflet give essential information to healthcare professionals and patients on how Teriflunomide should be used.

Important new concerns or changes to the current ones will be included in updates of Teriflunomide RMP.

I. The Medicine and What It is used for

Teriflunomide is authorised for the treatment of adult patients and paediatric patients aged 10 years and older with relapsing remitting multiple sclerosis (MS) (see SmPC for the full indication). It contains teriflunomide as the active substance and it is taken by mouth.

II. Risks Associated with the Medicine and Activities to Minimise or Further Characterise the Risks

Important risks of Teriflunomide, together with measures to minimise such risks are outlined below.

Measures to minimise the risks identified for medicinal products can be:

- Specific information, such as warnings, precautions, and advice on correct use, in the package leaflet and SmPC addressed to patients and healthcare professionals;
- Important advice on the medicine's packaging;
- The authorised pack size — the amount of medicine in a pack is chosen so to ensure that the medicine is used correctly;
- The medicine's legal status — the way a medicine is supplied to the patient (e.g. with or without prescription) can help to minimise its risks.

Together, these measures constitute *routine risk minimisation* measures.

In the case of Teriflunomide these measures are supplemented with *additional risk minimisation measures* mentioned under relevant important risks, below.

In addition to these measures, information about adverse reactions is collected continuously and regularly analysed, including PSUR assessment, so that immediate action can be taken as necessary. These measures constitute *routine pharmacovigilance activities*.

II.A List of Important Risks and Missing Information

Important risks of Teriflunomide are risks that need special risk management activities to further investigate or minimise the risk, so that the medicinal product can be safely taken. Important risks can be regarded as identified or potential. Identified risks are concerns for which there is sufficient proof of a link with the use of Teriflunomide. Potential risks are concerns for which an association with the use of this medicine is possible based on available data, but this association has not been established yet and needs further evaluation. Missing information refers to information on the safety of the medicinal product that is currently missing and needs to be collected (e.g. on the long-term use of the medicine).

Table 10: Summary of Safety Concerns

| List of important risks and missing information | |
|--|---|
| Important identified risks | <ul style="list-style-type: none"> • Hepatic effects • Hypertension • Haematologic effects • Infections • Acute pancreatitis |
| Important potential risks | <ul style="list-style-type: none"> • Teratogenicity • Serious opportunistic infections, including progressive multifocal leukoencephalopathy (PML) |
| Missing information | <ul style="list-style-type: none"> • None. |

II.B Summary of Important Risks

The safety information in the proposed Product Information is aligned to the reference medicinal product.

| Important identified risk: Hepatic effects | |
|--|--|
| Risk minimisation measures | <p><u>Routine risk minimisation measures:</u> SmPC sections 4.2, 4.3, 4.4, 4.8 and 5.2. Prescription only medicine. Prescription should be initiated and supervised by physicians experienced in the management of MS.</p> <p><u>Additional risk minimisation measures:</u> Educational Material for Health Care Professionals (HCP): HCP education/discussion guide; Educational Material for Patients: Patient education card.</p> |
| Important identified risk: Hypertension | |
| Risk minimisation measures | <p><u>Routine risk minimisation measures:</u> SmPC sections 4.4 and 4.8. Prescription only medicine. Prescription should be initiated and supervised by physicians experienced in the management of MS.</p> <p><u>Additional risk minimisation measures:</u> Educational Material for Health Care Professionals (HCP): HCP education/discussion guide; Educational Material for Patients: Patient education card.</p> |
| Important identified risk: Haematologic Effects | |
| Risk minimisation measures | <p><u>Routine risk minimisation measures:</u> SmPC sections 4.3, 4.4, 4.8 and 5.1. Prescription only medicine. Prescription should be initiated and supervised by physicians experienced in the management of MS.</p> <p><u>Additional risk minimisation measures:</u> Educational Material for Health Care Professionals (HCP): HCP education/discussion guide; Educational Material for Patients: Patient education card.</p> |

| Important identified risk: Infections | |
|---|--|
| Risk minimisation measures | <p><u>Routine risk minimisation measures:</u> SmPC sections 4.3, 4.4 and 4.8. Prescription only medicine. Prescription should be initiated and supervised by physicians experienced in the management of MS.</p> <p><u>Additional risk minimisation measures:</u> Educational Material for Health Care Professionals (HCP): HCP education/discussion guide; Educational Material for Patients: Patient education card.</p> |
| Important identified risk: Acute pancreatitis | |
| Risk minimisation measures | <p><u>Routine risk minimisation measures:</u> SmPC sections 4.4 and 4.8. Prescription only medicine. Prescription should be initiated and supervised by physicians experienced in the management of MS.</p> <p><u>Additional risk minimisation measures:</u> None</p> |
| Important potential risk: Teratogenicity | |
| Risk minimisation measures | <p><u>Routine risk minimisation measures:</u> SmPC sections 4.3, 4.6 and 5.3. Prescription only medicine. Prescription should be initiated and supervised by physicians experienced in the management of MS.</p> <p><u>Additional risk minimisation measures:</u> Educational Material for Health Care Professionals (HCP): HCP education/discussion guide; Educational Material for Patients: Patient education card.</p> |
| Important potential risk: Serious opportunistic infections, including progressive multifocal leukoencephalopathy (PML) | |
| Risk minimisation measures | <p><u>Routine risk minimisation measures:</u> SmPC sections 4.3, 4.4 and 4.8. Prescription only medicine. Prescription should be initiated and supervised by physicians experienced in the management of MS.</p> <p><u>Additional risk minimisation measures:</u> Educational Material for Health Care Professionals (HCP): HCP education/discussion guide; Educational Material for Patients: Patient education card.</p> |

II.C Post-Authorisation Development Plan

II.C.1 Studies Which Are Conditions of the Marketing Authorisation

There are no studies which are conditions of the marketing authorisation or specific obligation of Teriflunomide.

II.C.2 Other Studies in Post-Authorisation Development Plan

There are no studies required for Teva's Teriflunomide (Teriflunomid-ratiopharm).